

## **A INVESTIGATIONAL STUDY OF *ANTIBACTERIAL* ACTIVITIES OF *NIGELLA SATIVA* ON MASTITIS IN DAIRY CROSSBRED COWS**

**T. Monika, P. Sasikala., Dr. M. Vijaya Bhaskara Reddy\***

Dept. of Animal Nutrition, College of Veterinary University, Sri Venkateswara Veterinary  
University, Rajendra Nagar, India, +91-9052224389

Department of LPM, College of Veterinary University, Sri Venkateswara Veterinary University,  
Tirupati, India, +91-7396120530,+91-9705497313

---

### **Abstract**

In India, *Nigella Sativa* (*N. Sativa*) seeds were used in traditional medicine for the treatment of a variety of diseases in mammals and animals. The seed extracts and oil of this plant have shown various pharmacological properties including antimicrobial activities. The present study was aimed to investigate *in vitro* and *in vivo* antibacterial effects of methanol extract of the seeds against pathogenic bacteria causing mastitis in cows during the year 2010-11. *In vivo* studies in cows with mastitis were treated by local injection of different concentrations of methanol extract of the seeds into the infected breasts. *In vitro* experiments, the microorganisms were collected from the same infected breasts and used for the assessment of the antimicrobial effects of the extract by means of agar dilution and disk diffusion methods. The extract showed significant *in vitro* and *in vivo* inhibitory effects on causative organisms compared to standard drugs and also induced healing of the disease. This is the first experiment in crossbred cows in India, especially investigations on the antibacterial effects of *Nigella sativa*.

**Keywords:** Mastitis, *Nigella sativa*, Blackseed, Cross Bred Cows and Plant extracts.

---

### **Introduction**

Affections related to udder are to be given paramount importance as these affect economic conditions of farmers. Among different conditions, affecting mammary system udder

---

edema and mastitis is one of the important conditions, which clinically appears as an excessive accumulation of fluid in the interstitial spaces. It results in decreased milk production and makes difficulty in suckling and loss of productivity (V.B. Reddy *et al* 2013). *Nigella sativa* (blackseed) is common name in telugu is nalla jilakarra (నల్ల జీలకర్ర), an annual Ranunculaceae herbaceous plant growing to 30 cm and has an upright branching stem, fine deeply cut leaves, gray-blue flowers and toothed seedpots. The plant is native to Western Asia and the Mediterranean region (fig: 1). The seeds contain 40% fixed oil, saponins (melantin) and up to 1.4% volatile oil (Chevallier, 1996). Dioscorides, a Greek physician of the 1<sup>st</sup> century AD, recorded that black cumin seeds were taken to treat headaches, nasal catarrh, toothache, and intestinal worms (Chevallier, 1996). The seeds of *N. sativa* have been used traditionally for centuries in the Middle East, Northern Africa and India for the treatment of various diseases (Brutis and Bucar, 2000; Gilani et al., 2004). The plant extracts and essential oil showed a broad range of pharmacological effects such as antidiabetic (Farah et al., 2002; Benhaddou-Andaloussi et al., 2010), spasmolytic and bronchodilator (Gilani et al., 2001; Boskabady et al., 2008), hepatoprotective (Al-Ghamdi, 2003; Coban et al., 2010), analgesic and anti-inflammatory (Hajhashemi et al., 2004; Bashir and Qureshi, 2010), antitumor (Khan et al., 2003; Majdalawieh et al., 2010) and gastroprotective (El-Dakhakhny et al., 2000; Kanter et al., 2006) effects in various studies. The extracts also showed *in vitro* and *in vivo* antimicrobial (Mashhadian and Rakhshandeh, 2005; Salem et al., 2010), and anticestodal effects (Akhtar and Riffat, 1991). It is used traditionally in Iran as laxative, carminative and intestinal antiprotozoal drug (Amin, 1990). Mastitis is an inflammation of mammary gland regardless of the cause, with economical and health consequences. Many infective agents have been implicated as causes of mastitis; among the major pathogens are *S. aureus*, *E. coli*, streptococcus strains and *enterobacter aerogens* (Radostits et al., 1994). There is an increasing problem with antimicrobial drug resistance and so increasing demand for the development of new antimicrobial agents, especially in common veterinary and human infections. As there is no previous study on veterinary infections due to pathogenic microbes, the present study is initiated to investigate, *in vitro* and *in vivo* effects of methanol extract of blackseed on mastitis in crossbred cows.

## Materials and methods

*Nigella sativa* seeds were collected from local herbal drugs shop in tirupati. Chemicals Methanol and Streptomycin and penicillin G were purchased from (Hi Media (P) Ltd Mumbai).

### Reflux extraction

100 g of Blackseed powder were used with 600 ml of methanol with an extraction period of 12 hours. The extract was filtered using whatmann filter paper and the solvent was evaporated using rotary distillation apparatus. In order to obtain a completely dry extract, the resultant extract was transferred to glass dishes and was left in a 50°C oven for 24 hours. Then, it was stored at 4°C until assessment of antimicrobial activity.

### Clinical experiments

Treatment of mastitis in cows was carried out in Dairy experimental station, Dept. of LPM, CVSc, SVVU, Tirupati. Mastitis and the results of treatment were confirmed by a veterinary surgeon. Group A: 5 ml of 1% and 3% solutions of methanol extract of the seeds in liquid paraffin were injected into the four quarters (Teats) once daily for 5 consecutive days for the treatment of clinically-confirmed mastitis in crossbred cows. Group B: animals, treated with 10 ml of 6% solutions of the extract were injected twice daily at 12 hours interval. Negative control group: C treated with liquid paraffin only. Milk samples were collected from all animals before experimentation and 2 days after treatment and assayed for the presence of pathogenic bacteria. To avoid any contamination, sampling was carried out after careful washing of the breasts and disinfecting with povidine iodine. In order to calculate the reduction of colony count due to treatments, the following formula was used:

$$\% \text{ of reduction in colony count} = \frac{100 - \text{colony count after treatment}}{\text{Colony count before treatment}} \times 100$$

**In vitro microbiological assessments:** In vitro experiments were carried out in Group B only.

### Agar dilution method

Pre-defined amounts (Table 3) of the extracts were added to 100 ml of Muller- Hinton agar culture media (Merck) to obtain different concentrations in the media. Plates containing 15

ml of these media were incubated in 25°C for a period of 24 h to assure that they have not been contaminated during preparation. The microorganisms were cultured on these plates, incubated as mentioned above and the results were recorded and analysed for further studies.

### **Disk diffusion method**

Previously weighed paper disks were immersed for half an hour in the solutions of different concentrations of the extracts and were dried out under a laminar flow cabinet. By weighing the dried disks and comparing with before immersion weights, the amount of extracts in disks were calculated. Negative control disks were prepared using the solvent of the extracts in the same way. These (together with positive control disks, containing penicillin G and streptomycin) have been used for microbiological assay in plates containing the appropriate microorganisms.

### **Statistical analysis**

Paired t-test was used for the statistical analysis of the results of clinical experiments. Analysis of in vitro experiments was carried out using ANOVA and Tukey tests.

### **Results**

The results of the clinically collected samples of microorganism from the group A and B are shown in Table 1. In Group A, The results of colony count before and after once daily treatment of mastitis with 5 ml of 1% and 3% solutions of methanol extract of *N. sativa* seeds in paraffin are shown in Tables 2. The results indicate that the extract had a stronger effect against *S. aureus* than against group B *ι-hemolytic streptococci*. Where as in group B, The elevated results shows the no of colony count before and after twice daily treatment of mastitis with 10 ml of 6% solution of methanol extract of the seeds in paraffin are shown in Table 2. The results indicate that the extract had a strong effect against group D *streptococci*, *E. coli* and *S. aureus* in this concentration. The inhibitory effect was weaker against *E. coli* than against the other two organisms when compared with each other.

### **In vitro studies:**

As stated above, these experiments were carried out only on Group B samples; the antibacterial effects of methanol extract of *N. sativa* seeds on microorganisms isolated from the udder of the affected cows were investigated. Most common organisms identified in these experiments were *Enterococci* (group D *Streptococci*), *E. coli* and *S. aureus*, microbiological assessment was focused on these organisms. Table 1 shows the number of collected samples for the microorganisms.

Table: 1. Collected samples of milk from the affected cows.

Sample	No. Organism	No. of cases	
		Group A	Group B
1	<i>S. aureus</i>	15	5
2	Group B streptococci	12	0
3	<i>S. epidermis</i>	4	2
4	<i>E. coli</i>	3	15
5	<i>P. aeruginosa</i>	2	2
6	Group A streptococci	3	0
7	<i>Corinebacterium</i>	2	0
8	<i>Enterococci</i> (Group D streptococci)	0	40
9	<i>Klebsiella</i>	0	2
10	<i>Citrobacter</i>	0	1
11	Sterile	*	8
	Total	41	75

### **Agar dilution method**

The results of the current study shown in Table 3, the minimum inhibitory concentration (MIC) of the extract were 20 mg/ml (all samples of the organisms).

### **Disk diffusion method**

Table 4 shows the zones of inhibition for different concentrations of the extract against collected samples of *S. aureus*, *E. coli* and *Enterococci*. The extract did not show any effect against *E. coli* in this method. The amount of the extract in the disk for zone of inhibition of 10-20 mm for the collected samples of *S. aureus* and *Enterococci* were 1.5 mg and 5 mg,

respectively. The extract did not show any activity against *E. coli*. Zone of inhibition of more than 20 mm was formed only for collected strains of *CPSA* in 8 mg disks .



Fig: 1. *Nigella sativa* seeds and plant with flowers (ನಲ್ಲ ಜೀಲಕಠ)

Table 2. The *in vivo* antibacterial effect of methanol extract of *Nigella sativa* seeds against different infecting bacteria (Colony count in 1 ml of milk  $\times$  1000).

Organism	Collected from	Extract concentration	N	Before treatment	After treatment	Anova
<i>S. aureus</i>	Group A	1%	10	62 $\pm$ 10.8	34 $\pm$ 9.1	T= P
	Group A	3%	10	51 $\pm$ 11.9	17.3 $\pm$ 7.8	*p<0.05
<i>b-hemolytic streptococci</i>	Group B	1%	12	45 $\pm$ 8.3	42.7 $\pm$ 7.6	ns
	Group B	3%	11	38.2 $\pm$ 9.1	18.7 $\pm$ 6.8	*p<0.05
<i>Enterococci (Group D streptococci)</i>	Group B	6%	15	45.87 $\pm$ 8.98	10.7 $\pm$ 4.4	**p< 0.01
<i>E. coli</i>	Group B	6%	12	19.2 $\pm$ 8.4	0 $\pm$ 0	*p<0.05
<i>S. aureus</i>	Group B	6%	5	42 $\pm$ 12	0 $\pm$ 0	** p< 0.01

Data are presented as mean $\pm$  SEM.

Table 3. The *in vitro* antibacterial effect of methanol extract of *Nigella sativa* seeds against organisms collected in farm 2 in agar dilution method.

Extract concentration (mg/ml)	Enterococci Group-D Streptococci		E. Coli		S. Aureus	
	Growth %	Growth inhibition %	Growth %	Growth inhibition %	Growth %	Growth inhibition %
0	100	0	100	0	100	0
20	25	75	90	0	30	70
40	0	100	30	70	0	100
80	0	100	0	100	0	100
160	0	100	0	100	0	100

Table 4. Zones of inhibition produced by the extract against organisms collected in farm 2 in disk diffusion method.

Organism	Penicillin G	Dose	N	Zone of inhibition
<i>Enterococci</i>	Penicillin G	(10 mg/disk)	21	13.05 ± 1.39
	Streptomycin	(10 mg/disk)	21	10.04 ± 1.9
<i>S. Aureus</i>	Penicillin G	(10 mg/disk)	5	9.8 ± 4.3
	Streptomycin	(10 mg/disk)	5	21 ± 2.32
<i>Enterococci</i>	Extract	(5 mg/disk)	21	11.28 ± 1.28
		(8 mg/disk)	21	13.23 ± 1.27
<i>S. Aureus</i>	Extract	(1.5 mg/disk)	5	10.4 ± 2.6
		(4 mg/disk)	5	17 ± 1.34
		(8 mg/disk)	5	20.2 ± 1.32

There were no significant responses with concentrations below 4 mg/disk and 2 mg/disk against *Enterococci* or *S. aureus* respectively. Also, methanol did not produce any zone of inhibition. Data presented as mean  $\pm$  SEM.

## Discussion

Statistical analysis of the results showed that the methanol extract of *N. sativa* seeds had significant *in vivo* antimicrobial activity in Group A and B experiments and these effects were dose-dependent. Treatment of clinically-confirmed mastitis with injection of 10 ml of 6% paraffin solution of the extract twice daily for 5 days induced complete response against *E. coli* and *S. aureus* and an excellent response against *Enterococci* (group D *Streptococci*) in group b (Table 2). Injection of 5 ml of 1% and 3% paraffin solution of the extract once daily also produced significant antimicrobial activity against *S. aureus* in group B but only 3% solution produced significant activity against group B  $\beta$ -hemolytic *Streptococci* in B group (Table 2). It is obvious that the use of higher dose of the extract in B group (10 ml of 6% solution twice daily compared with 5 ml of 1% and 3% twice daily in A group) produced stronger effects against the organisms, as shown for *S. aureus* in Table 2. Results of *in vitro* experiments also showed significant activity of the extract against *S. aureus* and *Enterococci* in all methods and against *E. coli* in agar dilution method only (Table 3). Hanafy and Hatem (1991) also observed antimicrobial activity of diethyl ether extract of the plant against *S. aureus*, *P. aeruginosa*, *E. coli* and *C. albicans*. Our results did not show any activity against *E. coli* in disk diffusion method. Due to methanol extract the results affected some extent. The amount of ingredients of the same plant can be affected by the area and the season of collection. Sometimes, resistant strains of some organisms exist in animal setting compared with standard or human ones. This extract was active against *E. coli* in our clinical experiments. Finally, *in vivo* effects of antimicrobial agents may be different from their *in vitro* effects, because of the effects of immune system. It seems that immunological and environmental factors affect the activity of the extract against *E. coli*. *N. sativa* seed extracts also showed anticestodal (Akhtar and Riffat, 1991). Finally, our results are in agreement with others who showed that *N. sativa* extracts produce antimicrobial activity against a broad range of microbes and especially against multiple-antibiotic resistant bacteria (Morsi, 2000). This is the first ever report on *in vivo* study of antibacterial effects of *N. sativa* seed

extracts against a common and important disease of cattle. Considers that the wide margin of safety of the extracts of the seeds (Gilani et al., 2004; Vahdati- Mashhadian et al., 2005).

The mechanism of action of antimicrobial effects of the extracts is not clear but their broad spectrum of activity implies that they are affecting basic and common key processes in the organisms. They may be of promising veterinary uses in the future. Further studies on the activity-directed fractionation for the isolation of respective pure compounds may result in interesting outcomes.

**ACKNOWLEDGEMENT:** It is with most profound feelings of respect, sincerity and high regards that I express my indebtedness and deep sense of gratitude to my venerable brother, Pioneer and teacher M. Chandrasekhara Reddy & family K. Bharani, Master M. Druvin Reddy, SWE, Ohio, USA, for his meticulous guidance, Stimulating discussion, Awe inspiring Encouragement and suggestions to complete this work with confidence. I will ever remain grateful to him for his inspiring guidance, wise counsel unfailing attention. He has been a source of inspiration and confidence in my research work.

**REFERENCES:**

- Akhtar MS, Riffat S. 1991. Field trial of *Saussurea lappa* roots against nematodes and *Nigella sativa* against cestodes in children. *J Pak Med Assoc*, 41(8):185-7.
- Al-Ghamdi MS. 2003. Protective effect of *Nigella sativa* seeds against carbontetrachloride-induced liver damage. *Am J Chin Med*, 31:721-8.
- Amin G. 1990. *Traditional herbal drugs in Iran*. Iranian ministry of health publications, Tehran, PP: 118-119.
- Bashir MU, Qureshi HJ. 2010. Analgesic effect of *Nigella sativa* seeds extract on experimentally induced pain in albino mice. *J Coll Physicians Surg Pak*, 20:464-7.
- Benhaddou-Andaloussi A, Martineau LC, Vallerand D, Haddad Y, Afshar A, Settaf A, Haddad PS. 2010. Multiple molecular targets underlie the antidiabetic effect of *Nigella sativa* seed extract in skeletal muscle, adipocyte and liver cells. *Diabetes Obes Metab*, 12:148-57.
- Boskabady MH, Keyhanmanesh R, Saadatloo MA. 2008. Relaxant effects of different fractions from *Nigella sativa* L. on guinea pig tracheal chains and its possible mechanism(s). *Indian J Exp Biol*, 46:805-10.

- Brutis M, Bucar F. 2000. Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res*, 14:323-8.
- Chevallier A. 1996. The encyclopedia of medicinal plants, Dorling Kindersley publishers, London, P: 237.
- Coban S, Yildiz F, Terzi A, Al B, Aksoy N, Bitiren M, Celik H. 2010. The effects of *Nigella sativa* on bile duct ligation induced liver injury in rats. *Cell Biochem Funct* 28: 83-8.
- El-Dakhakhny M, Barakat M, El-Halim MA, Aly SM. 2000. Effects of *Nigella sativa* oil on gastric secretion and ethanol induced ulcer in rats. *J Ethnopharmacol*, 72(1-2):299- 304.
- Fararh KM, Atoji Y, Shimizu Y, Takewaki T. 2002. Insulinotropic properties of *Nigella sativa* oil in Streptozotocin plus Nicotinamide diabetic hamster. *Res Vet Sci* 73(3):279-82.
- Gilani AH, Aziz N, Khurram IM, Chaudhary KS, Iqbal A. 2001. Bronchodilator, spasmolytic and calcium antagonist activities of *Nigella sativa* seeds (Kalonji): a traditional herbal product with multiple medicinal uses. *J Pak Med Assoc* 51(3):115- 20.
- Gilani AH, Jabeen Q, Khan MAU. 2004. A review of medicinal uses and pharmacological activities of *Nigella sativa*. *Pak J Biol Sci* 7:441-51.
- Hajhashemi V, Ghannadi A, Jafarabadi H. 2004. Black cumin seed essential oil, as a potent analgesic and antiinflammatory drug. *Phytother Res* 18:195-9.
- Hanafy MS, Hatem ME. 1991. Studies on the antimicrobial activity of *Nigella sativa* seed (black cumin). *J Ethnopharmacol*, 34:275-8.
- Kanter M, Coskun O, Uysal H. 2006. The antioxidative and antihistaminic effect of *Nigella sativa* and its major constituent, thymoquinone, on ethanol-induced gastric mucosal damage. *Arch Toxicol* 80:217-24.
- Majdalawieh AF, Hmaidan R, Carr RI. 2010. *Nigella sativa* modulates splenocyte proliferation, Th1/Th2 cytokine profile, macrophage function and NK anti-tumor activity. *J Ethnopharmacol* 131:268-75.
- Mashhadian NV, Rakhshandeh H. 2005. Antibacterial and antifungal effects of *Nigella sativa* extracts against *S. aureus*, *P. aeruginosa* and *C. albicans*. *Pak J Med Sci* 21:47-52.
- Morsi NM. 2000. Antimicrobial effect of crude extracts of *Nigella sativa* on multiple antibiotics-resistant bacteria. *Acta Microbiol Pol* 49:63-74.

- Radostits OM, Blood DC, Gay CC. 1994. Veterinary medicine. London, W. B. Saunders, PP: 563-4.
- Salem EM, Yar T, Bamosa AO, Al-Quorain A, Yasawy MI, Alsulaiman RM, Randhawa MA. 2010. Comparative study of *Nigella Sativa* and triple therapy in eradication of Helicobacter Pylori in patients with nonulcer dyspepsia. Saudi J Gastroenterol, 16:207-14.
- Vahdati-Mashhadian N, Rakhshandeh H, Omid A. 2005. An investigation on LD50 and subacute hepatic toxicity of *Nigella sativa* seed extracts in mice. Pharmazie, 60:544-7.
- Vijaya Bhaskara Reddy. M, Sasikala. P and Sreedhar. S. 2013. Extensive Udder Edema Management In Jersey X Sahiwal Crossbred Cows Under Field Conditions. Int, Jr. Adv. Sci. Tech. Res. 3: 549-553.